

# Michigan Collaborative Standardization of Compounded Oral Liquids

---

<b>Drug name:</b>	<b>Buspirone</b>	<b>Dosage Form:</b>	Suspension
<b>Concentration:</b>	2.5 mg/ml	<b>Shelf Life:</b>	30 days
<b>Route:</b>	Oral	<b>Storage:</b>	Refrigerate or Room Temp
<b>Volume:</b>	100 ml	<b>Auxiliary Labeling:</b>	Shake Well

Ingredients	QS	Quantity	Units
Buspirone 10 mg tablets		25	tablets
Ora-Blend or Ora-Plus/Ora-Sweet (1:1)	Y	100	mLs

## Directions:

1. Crush tablets in a mortar and pestle and triturate to a fine powder.
2. Add a minimal amount of vehicle to cover the tables and allow them to soak.
3. Mix with pestle into a smooth suspension.
4. Continue adding vehicle in geometric portions, mixing well.
5. Transfer to graduate.
6. Rinse mortar and pestle with vehicle.
7. QS to final volume with vehicle. Stir well.

## Notes:

- 

## References:

- Edwards DF, Chugani DC, Chugani HT et al. J Clin Pharmacology. 2006; 46(5): 508-14.

Reviewed: April 2013; SC\JR

The information provided on this site is only to be used for educational and informational purposes only. It may not be relied upon to direct or provide care to a patient, or to prepare treatments for a patient. This material is not intended as a substitute for retaining and consulting appropriate medical and health care professionals, who remain solely responsible to provide and direct patient care. Be aware that the information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professionals or pharmacists, and may require governmental approvals. The owner and provider of this site disclaims and assumes no liability for any injury and/or damage to persons from use of any information contained on this site.

---