

Michigan Collaborative Standardization of Compounded Oral Liquids

Drug name: Desmopressin

Concentration: 0.033 mg/mL

Route: Oral

Volume: 7.5 mL

Dosage Form: Solution

Shelf Life: 30 days

Storage: Refrigerate

Auxiliary Labeling: Refrigerate

Ingredients	QS	Quantity	Units
Desmopressin acetate solution 0.1 mg/mL		2.5	mL
NaCl 0.9% injection, USP (nonbacteriostatic)		5	mL

Directions:

1. Add NaCl 0.9% injection to desmopressin solution bottle using a sterile syringe and needle
2. Shake
3. Pull doses from bottle using an IV syringe and transfer to oral syringes

Notes:

- Must be prepared in IV Hood

References:

- Nahata MC, Pai VB. *Pediatric Drug Formulations*. 6th ed. Cincinnati, OH: Harvey Whitney Book; 2011.

Reviewed: April 2013; ME/JR

The information provided on this site is only to be used for educational and informational purposes only. It may not be relied upon to direct or provide care to a patient, or to prepare treatments for a patient. This material is not intended as a substitute for retaining and consulting appropriate medical and health care professionals, who remain solely responsible to provide and direct patient care. Be aware that the information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professionals or pharmacists, and may require governmental approvals. The owner and provider of this site disclaims and assumes no liability for any injury and/or damage to persons from use of any information contained on this site.

