

# Michigan Collaborative Standardization of Compounded Oral Liquids

---

<b>Drug name:</b>	<b>Tretinoin</b>	<b>Dosage Form:</b>	Suspension
<b>Concentration:</b>	10 mg/ml	<b>Shelf Life:</b>	30 days
<b>Route:</b>	Oral	<b>Storage:</b>	Room temp
<b>Volume:</b>	100 ml	<b>Auxiliary Labeling:</b>	Shake Well; Protect from Light; Antineoplastic agent

Ingredients	QS	Quantity	Units
Tretinoin		1	gm
Butylated hydroxyanisole		0.02	gm
Edetate disodium dehydrate		0.1	gm
Soybean oil	Y	100	mL

## Directions:

1. Calculate the required quantity of each ingredient for the total amount to be prepared
2. Weigh and/or measure each ingredient.
3. Add a portion of the soybean oil to the powders and mix well using a disposable mortar and pestle or by the syringe method.
4. Add mixture to calibrated glass amber bottle and add soybean oil to bring the mixture to desired volume.
5. Cap and shake well.

## Notes:

- Caution: Antineoplastic agent - preparation, including weighing of materials, requires hazardous chemical precautions.

## References:

- McElhiney LF. Inter J Pharm Compounding. 2010; 14(2): 118-23.

Reviewed: April 2013; SC\ME

The information provided on this site is only to be used for educational and informational purposes only. It may not be relied upon to direct or provide care to a patient, or to prepare treatments for a patient. This material is not intended as a substitute for retaining and consulting appropriate medical and health care professionals, who remain solely responsible to provide and direct patient care. Be aware that the information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professionals or pharmacists, and may require governmental approvals. The owner and provider of this site disclaims and assumes no liability for any injury and/or damage to persons from use of any information contained on this site.

---